



### What is a Cervical Incompetence?

The cervix normally gradually weakens from the pregnancy hormones that stimulate it to dilate and allow the fetus to pass. It can sometimes be the case it weakens a little earlier than normal and causes a Cervical Incompetence. The Tanner Clinic Cervical Incompetence Programme is a new [open access programme at Tanner Clinic aimed at the UK and Irish](#).

We have a team of experts who specialise in Cervical Incompetence (CIC) care. We are also members of the British Fertility Society, which means we have a strong reputation and a track record with a team of us.

### What are the symptoms of CIC/ICP?

The symptoms of pregnancy loss that occur and you do not have a history of miscarriages or stillbirths. Symptoms like heavy and watery discharge from a woman with CIC/ICP. The TIC will agree [a treatment](#) if more evidence is available.

### What are the chances of CIC/ICP occurring with a subsequent pregnancy?

There is a 50% chance of a subsequent pregnancy loss or a pregnancy party. Find the [TIC](#) and the [TIC programme at Tanner](#) programme website.

### What are the most reliable ways to measure your risk from CIC/ICP?

Additional blood and urine testing will allow us to find out if you have a CIC/ICP and assess your pregnancy. The TIC is a comprehensive programme [TIC/ICP](#) and a great resource for when you are not pregnant. Contact us for more information. Also, be prepared to pay us.