

When the plant is in the vegetative phase, when given a high concentration of auxin, it will produce an abscisic acid-like compound. This compound will be transported through the plant and will cause the plant to stop growing and to produce a dormancy-like state. This is the mechanism by which the plant can survive in a dormant state.

**[[Auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa.]]**

**Question on top - 10/10/2020, 10/10/2020**

### **10/10/2020 change summary:**

1. The first is a very general one. It has been shown that auxin is a key hormone in the growth of plants. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.
2. **10/10/2020 change summary:**
  1. The first is a very general one. It has been shown that auxin is a key hormone in the growth of plants. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.
  2. The second is a very specific one. It has been shown that auxin is a key hormone in the growth of plants. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.
  3. The third is a very specific one. It has been shown that auxin is a key hormone in the growth of plants. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.

### **10/10/2020 change summary:**

1. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
2. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
3. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
4. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
5. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
6. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
7. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
8. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
9. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**
10. **Change in auxin transport - In the plant, auxin is transported from the shoot to the root and vice versa. It is a hormone that is produced in the shoot and is transported to the root. It is a hormone that is produced in the shoot and is transported to the root.**